

Amprolium

COMPOSITION

Each 30 g contains:

Amprolium hydrochloride 6 g

INDICATION

For treatment & prevention of coccidiosis in replacement chicks, layers, breeders, broilers & turkeys

DOSAGE & ADMINISTRATION

Dissolve Amprolium soluble powder in drinking water as follows:

For prevention : 30 g in 50 litres of water

For mild infection : 30 g in 25 litres of water

For disease outbreaks : 60 g in 25 litres of water

or as recommended by Veterinarian

Medicated water should be given for 5-7 days. No other source of water should be available to the birds, other than the medicated water. Once the medicated water is completely consumed, replenish with fresh non medicated water.

Amprolium soluble powder is a safe anticoccidial having no residues & is compatible with vitamins (except vitamin B₁), antibiotics & minerals.

PRESENTATION

Sachet of 150 g • Bag of 1 kg



ANTICOCCIDIALS

SUPERCOX®

COMPOSITION

Each 100 g contains	
Sulphaquinaxaline	18.7 g
Diaveridine	3.3 g
Excipients	q.s.

INDICATIONS

For early & effective control of intestinal & caecal coccidiosis in poultry caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. praecox* & *E. mitis*.

Fowl cholera caused by *Pasteurella multocida*. Fowl typhoid caused by *Salmonella gallinarum*.

DOSAGE & ADMINISTRATION

10 g / 10 litre water or 100 g in 50 kg feed for 3 days

Plain water / Feed for next 2 days

10 g / 20 litre water or 100 g in 100 kg feed for next 3 days

or as recommended by Veterinarian

PRESENTATION

Sachet of 100 g



2



NUTRACEUTICALS

COMPOSITION

Each kg contains:	
Vitamin A	26,43,200 I.U.
Vitamin D ₃	4,00,000 I.U.
Vitamin E	1,111 I.U.
Vitamin C	5,00,000 I.U.
Vitamin K ₃	1.104 g
Riboflavin (B ₂)	2.864 g
Niacin (B ₃)	13.52 g
Ca D Pantothenate (B ₅)	4.408 g
Pyridoxine (B ₆)	552 mg
Cyanocobalamin (B ₁₂)	2.20 mg
MSM	10.00 g
Betaine	50 g
L-Selenomethionine	100 mg

USAGE

Various type of stress conditions in poultry like handling, brooding, vaccination, debeaking etc.

FEEDING DIRECTION : 1g / litre of drinking water

PRESENTATION : 1kg



3



NUTRACEUTICALS

Caritol™

COMPOSITION

Each ml contains	
Sorbitol	200 mg
D-L Methionine	10 mg
Choline chloride	18.75 mg
Magnesium sulphate	10 mg
L-Carnitine hydrochloride	25 mg
Aqueous base	q.s.

USAGE

- Elimination of toxins
- Support in fatty liver condition & other liver failures
- Support during or after antibiotic or antiparasitic treatment
- Helps in improvement of general health condition & production

FEEDING DIRECTION

Chicks : 5ml / 100 chicks
Growers : 10 ml / 100 birds
Adult Birds : 20 ml / 100 birds
To be administered through
drinking water daily for 5-7 days or
as advised by the Veterinarian



PRESENTATION : 1 & 5 Ltr.



NUTRACEUTICALS

arB Ce™

COMPOSITION

Liquid	
Each 10 ml contains	
Ferric ammonium citrate	1430 mg
(equivalent to 300 mg of Fe)	
Chelated Copper Sulphate	100 mg
Cobalt chloride	1.5 mg
Cholecalciferol	75 mg
Biotin	275 mcg
Folic Acid	1.5 mg
Cyanocobalamin	150 mcg
Excipients	q.s.

Feed Supplement

Each kg contains	
Chelated Iron55 gm
Chelated Copper Sulphate18 gm
Cobalt Chloride027 gm
Cholecalciferol	900000 IU
Nicotinamide	15 gm
Biotin	49500 mcg
Folic Acid	0.27 gm
Cyanocobalamin	27000 mcg
Excipients	q.s.



NUTRACEUTICALS



USAGE

To support in anemic conditions & to improve general health conditions

FEEDING DIRECTION

Broiler, Layer, Breeder Chicks :

10-15 ml/1000 chicks

Growers : 10 ml/100 birds

Layer, Broiler Adult : 20 ml/100 birds

Breeder Adult : 30 ml/100 birds

PRESENTATION

1ltr, 2 ltr, 20 Ltr. & 1 kg



ANTIBIOTICS

COMPOSITION

Each ml of Meriquin liquid contains
Enrofloxacin 100 mg

INDICATIONS

For treatment & prevention of infectious diseases caused by Gram Positive & Gram Negative bacteria, Mycoplasma spp. including mixed infections viz. C.C.R.D. Coli-septicaemia, Pasteurellosis, Coryza (Haemophilus spp.), Salmonella gallinarum & other Salmonella spp.

DOSAGE & ADMINISTRATION

For treatment & prophylaxis, administer 10 mg/kg B.wt. in drinking water for 3-5 days or as recommended by Veterinarian

PRESENTATION

Bottle of 100 ml & 500 ml

Jar of 5 liters



ANTIBIOTICS

Meriflox®

COMPOSITION

Liquid Levofloxacin Hemihydrate 10% w/v oral solution

INDICATIONS

For the control & treatment of infections in poultry like E.Coli infections, Salmonellosis, Fowl Cholera, Infectious Coryza, Dermatitis, CRD, CCRD etc.

Also controls other infections caused by Gram +ve & Gram -ve micro-organisms.

DOSAGE & ADMINISTRATION

- 1 ml / 10 kg BW
- To be administered continuously for a period of 3-5 days once daily in drinking water depending up on the severity of infection or as recommended by the Veterinarian

PRESENTATION

100 ml & 1 ltr., 5 ltr.



ANTIBIOTICS

Meriflox® BH

COMPOSITION

Each ml contains :
Levofloxacin hemihydrate I.P.
eq. to Levofloxacin 100 mg
Bromhexine HCl I.P. 7.5 mg

INDICATIONS

For the control & treatment of severe respiratory infections of poultry like CRD, CCRD, Infectious Coryza & Fowl Cholera

DOSAGE & ADMINISTRATION

- 1 ml / 10 kg BW
- To be administered continuously for a period of 3-5 days once daily in drinking water depending up on the severity of infection or as recommended by the Veterinarian

PRESENTATION

250 ml & 1ltr., 5 ltr.



TOXIN BINDER

TOXORID-BIO™

COMPOSITION

MOS
HSCAS
Organic acid
Activated Charcoal
Oxine Copper
Selenium

USAGE

For effective control of mycotoxins in feed

FEEDING DIRECTION

1 kg / ton of feed

PRESERNTATION

25 kg Polybag



10

ENZYME & PROBIOTICS

COMPOSITION

Each Kg contains	
Enzymes	
Cellulase	180,000,000 U
Amylase	125,000 U
Xylanase	1,800,000 U
Phytase	200,000 U
Protease	16,000 U
Lipase	40,000 U
Pectinase	7000 U

Probiotics

Lactobacillus acidophilus	1 billion CFU
Saccharomyces boulardii	1 billion CFU
Bacillus subtilis natto	1 billion CFU
Bacillus licheniformis	1 billion CFU

USAGE : For better nutrient digestion, utilization, gut health improvement & cost economization of feed formulation

FEEDING DIRECTION : 350 g per Tonne of Feed

PRESERNTATION : 25 Kg Polybag (5 x 5 kg) packets



11



ENZYME & PROBIOTICS

BIOVET-YC®

COMPOSITION:

Each kg of Biovet YC provides	
Saccharomyces boulardii	30,000 mio.c.f.u.
Lactobacillus acidophilus	45,000 mio.c.f.u.
Live yeast culture of	
Saccharomyces cerevisiae sc-47	3,00,000 mio. c.f.u.
Alpha amylase	5 g
Sea weed extract	100 g
Excipients	q.s.

USAGE:

- To prevent digestive upsets, diarrhoea etc.
- To exclude harmful pathogens from gut
- Improves Intestinal ecology

FEEDING DIRECTION:

Chick, grower & broiler : 500 g/tonne of feed
Layer & breeder : 1 kg/tonne of feed

PRESENTATION:

Poly pack of 1 kg



12



VITAMIN PREMIXES

Meriplex-K SS

COMPOSITION:

Each gm contains	
Vitamin A	40,000 I.U.
Vitamin D ₃	10,000 I.U.
Vitamin B ₁	3.2 mg
Vitamin B ₂	20 mg
Vitamin B ₆	6.4 mg
Vitamin B ₁₂	82 mcg
Niacin	48 mg
Calcium pantothenate (mg)	32 mg
Vitamin K ₃	4 mg
Vitamin E	32 mg
Folic acid	3.2 mg
Carriers	q.s.

USAGE:

Vitamin supplement to improve performance

FEEDING DIRECTION:

Mix 250 g / tonne of feed

PRESENTATION:

25 x 1 kg



13



VITAMIN PREMIXES

Merivite® 100

COMPOSITION

Each kg of Merivite 100 contains
Vitamin B₁₂ 100 mg

USAGE

Vitamin supplement to improve performance

FEEDING DIRECTION

Layer : 100 g per tonne of feed
Broiler/replacement : 220 g per tonne of feed
Breeder : 150 g per tonne of feed

PRESENTATION

Poly pack of 25 kg (5 x 5 kg)



14



OTHER

COMPOSITION

Each kg contains :
Erythromycin Thiocyanate 100 gm

USAGE

For effective mycoplasma control
Effective against CRD as well as other bacterial infection in upper respiratory tract

RECOMMENDED USAGE

For Commercial Broiler & Layers

- Mild infections : 200 gm/ton of feed
- Moderate infections : 350-500 gm/ton of feed
- High challenge areas : 1 kg/ton of feed

For Breeders : 2 to 3 kg/ton of feed

PRESENTATION

1 kg x 25 units in HDPE Drum



15

OTHER

RECYMETH®

COMPOSITION:

Methyl donors & methyl group recycling agents are easily bio available with organic sulphur base, fortified with short chain fatty acids, chelated minerals & Amino nitrogen

USAGE:

Alternative to Synthetic Methionine

FEEDING DIRECTION:

Mix with feed on equal weight basis of Synthetic Methionine

PRESENTATION:

25 kg poly bag (5 x 5 kg bag)



16

DISINFECTANT

ResimaxTM 15-10

COMPOSITION:

Glutaraldehyde	15% w/v
Benzalkonium Chloride	10% w/v
Pine Oil	q.s.
Terpineol	q.s.
Colour Food Grade - Tartrazine + Sunset Yellow (-)	
Purified Water	q.s.

DIRECTION FOR USE:

Completely safe for spraying in presence of birds 7 ml/ltr. 1 ltr. of dilute solution per 300 sq. ft.

Usage	Dilution	Application
Routine disinfection	7 ml per liter	300 ml of dilute solution per sq meter area
Surface disinfection	7 ml per liter	300 ml of dilute solution per sq meter area
Fogging	7 ml per liter	1 liter of dilute solution per 100 cubic meter
Foot dip / wheel wash	10 ml per liter	replace twice a week

PRESENTATION: 250 ml, 1 Ltr. & 5 Ltr.



17

